

Note to Readers: If you need assistance accessing items in this Supplemental Material, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Table of Contents for Supplemental Material

Ambient Air Pollution and Newborn Size and Adiposity at Birth: Differences by Maternal Ethnicity (the Born in Bradford Study Cohort)

Anna Schembari, Kees de Hoogh, Marie Pedersen, Payam Dadvand, David Martinez, Gerard
Hoek, Emily S. Petherick, John Wright, and Mark J. Nieuwenhuijsen

Table S1. Characteristics and exposure distributions for the full population and by maternal ethnicity.

Table S2. Pearson correlations between trimester specific exposure and the full pregnancy for each air pollutant.

Tables S3. Pearson correlations between air pollutant exposures for the full pregnancy.

Table S4. Adjusted models coefficients (95% CI) for air pollution exposure during 1st and 2nd trimester of pregnancy with birth weight (g) head circumference (cm), triceps and subscapular skinfolds thickness (mm).

Table S5. Term babies (N=8,586) - Adjusted models coefficients (95% CI) for air pollution exposure and birth weight (g) head circumference (cm), triceps and subscapular skinfolds thickness (mm).

Table S6. Mothers who don't change residence during pregnancy (N=8,521) - Adjusted models coefficients (95% CI) for ambient air pollution exposure and birth weight (g), head circumference (cm), triceps and subscapular skinfolds thickness (mm).

Table S7. Unemployed (n=4,436) vs. employed (n=3,201) mothers - Adjusted models coefficients (95% CI) for air pollution exposure and birth weight (g) head circumference (cm) triceps and subscapular skinfolds thickness (mm).

Table S8. Non-smokers (N=7,005) vs smoker (N=1,645) mothers - Adjusted models coefficients (95% CI) for air pollution exposure and birth weight (g), head circumference (cm), triceps and subscapular skinfolds thickness (mm).